



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 557 270 B1**

12

EUROPEAN PATENT SPECIFICATION

- 40 Date of publication of patent specification: 31.05.95
- 51 Int. Cl. A61K 39/395, C07K 16/18, //C12P21/00
- 21 Application number: 88907980.2
- 22 Date of filing: 15.08.88
- 66 International application number: PCT/US88/02760
- 67 International publication number: WO 89/01343 (23.02.89 89/05)

CATIONIZED ANTIBODIES FOR DELIVERY THROUGH THE BLOOD-BRAIN BARRIER.

- 30 Priority: 17.08.87 US 85627
- 43 Date of publication of application: 01.09.93 Bulletin 93/35
- 46 Publication of the grant of the patent: 31.05.95 Bulletin 95/22
- 64 Designated Contracting States: AT BE CH DE FR GB IT LI LU NL SE
- 56 References cited: US-A- 4 118 379

PHYSIOLOGICAL REVIEWS, vol. 22, 1942, pp. 125-145&NUM;

JOURNAL OF IMMUNOLOGY, vol. 133, no. 6, December 1984, American Association of Immunologists, US; B.H. HAHN et al., pp. 3015-3019&NUM;

CHEMICAL ABSTRACTS, vol. 107, 1987, Columbus, OH (US); W.M. PARDRIDGE et al., p. 524, no. 37764f&NUM;

- 73 Proprietor: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA
2199 Addison Street
Berkeley,
California 94720 (US)
- 72 Inventor: PARDRIDGE, William, M.
1180 Tellem Drive
Pacific Palisades, CA 90272 (US)
Inventor: SCHIMMEL, Paul, R.
31 Solomon Pierce Road
Lexington, MA 02173 (US)
- 74 Representative: Dealtry, Brian et al
Eric Potter & Clarkson
St. Mary's Court
St. Mary's Gate
Nottingham NG1 1LE (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Rank: Xerox (UK) Business Services
10/3.09/3.3.2

EP 0 557 270 B1

1

EP 0 557 270 B1

2

Description**Background of the Invention**

The present invention relates generally to the use of antibodies for treatment and diagnosis of neurological diseases. More particularly, the present invention deals with the modification of antibodies so that they can be delivered through the blood-brain barrier by transcytosis.

Antibodies in general, and especially monoclonal antibodies, are widely used in diagnostic tests as a means for detecting the presence of specific antigens. Enzyme linked immunoassay and radioimmunoassay are common diagnostic techniques which utilize antibodies and detect antigens in vivo. Antigens may also be detected in vitro by administering radiolabelled antibodies to a living subject followed by external detection of radiolabelled antibody sequestered by a particular organ bearing the respective antigen. Antibodies have also been used widely in the treatment of viral infections and other diseases. However, the use of antibodies in either the treatment or diagnosis of neurological diseases has been very limited because most antibodies are not capable of traversing the blood-brain barrier (BBB) and entering the brain.

The vertebrate brain has a unique capillary system which is unlike that in any other organ in the body. The unique capillary system has morphologic characteristics which make up the blood-brain barrier. The blood-brain barrier acts as a system wide cellular membrane which separates the brain interstitial space from the blood. The unique morphologic characteristics of the brain capillaries which make up the blood-brain barrier are: (a) epithelial-like high resistance tight junctions which literally cement all endothelia of brain capillaries together, and (b) scanty pinocytosis or transendothelial channels, which are abundant in endothelia of peripheral organs. Due to the unique characteristics of the blood-brain barrier, antibodies that readily gain access to other tissues in the body are barred from entry into the brain or their rates of entry are very low.

US-A-4118319 (Schmidtberger) discloses the covalent modification of carboxyl groups on immune globulins with primary amines.

Few strategies have been developed for introducing these antibodies into the brain which otherwise would not cross the blood-brain barrier. The most commonly used strategy involves an invasive procedure where the antibody is delivered directly into the brain. The most common procedure is the implantation of a catheter into the ventricular system to bypass the blood-brain barrier and deliver the antibody directly to the brain. Such

a procedure has been used in treating echovirus encephalitis (Erlendsson et al., Successful Reversal of Echovirus Encephalitis In X-linked Hypogammablobulinemia by Intraventricular Administration of Immunoglobulin 1985. New England Journal of Medicine. Vol. 312, No. 6. pages 351-353).

Although invasive procedures, such as the one described above, for the direct delivery of antibodies to the brain ventricles have experienced some success, they are not entirely satisfactory because they do not deliver the antibodies to the structures deep within the brain. Further, the invasive procedures are potentially harmful to the patient. Accordingly, there presently is a need to provide an improved method for delivering antibodies across the blood-brain barrier and into the brain for both diagnostic and therapeutic purposes.

Summary of the Invention

In accordance with the present invention, a method is provided for introducing antibodies directly across the blood-brain barrier by transcytosis. The invention is based upon the discovery that cationized antibodies cross the blood-brain barrier at much higher rates than normal non-cationized antibodies.

The effectiveness of antibodies for both neurodiagnostic and neuropharmaceutical purposes is increased by cationizing the antibodies to provide cationized antibodies having an isoelectric point (pI) of between about 8.0 to 11.0. These highly basic antibodies cross the blood-brain barrier at rates which are much higher than the transcytosis rates for normal acid and neutral antibodies which typically have isoelectric points in the range of 5 to 6. This provides an effective means for delivering antibodies into the brain by transcytosis rather than by the previously used direct invasive methods.

The cationized antibodies in accordance with the present invention are prepared by treating a given monoclonal or polyclonal antibody with a cationization agent such as hexamethylenediamine. The cationization agent replaces surface carboxyl groups on the antibody with a more basic group, such as a primary amine group in the case of hexamethylenediamine and related amine compounds. The amount of cationization agent and reaction conditions are controlled so that the resulting cationized antibody has an isoelectric point of between about 8.0 to 11.0 and preferably between about 8.0 to 9.0.

As one feature of the present invention, the immunoreactivity of the antibodies is preserved during cationization by first reacting the antibody with an excess of a corresponding antigen to block the immunoreactive sites on the antibody. These

2

3

EP 0 557 270 B1

4

blocked immunoreactive sites are unreactive during the subsequent cationization steps. The antigens are then decoupled from the cationized antibodies after the cationization step to thereby reactivate the blocked immunoreactive sites.

The cationization of antibodies in accordance with the present invention is useful whenever it is necessary to introduce an antibody into the brain. Both neurodiagnostic and neurotherapeutic uses for antibodies is contemplated. Particular diagnostic uses include diagnosis of Alzheimer's disease, brain tumors or any other diagnostic use where a labeled or tagged antibody is introduced into the brain for reaction with and detection of specific antigens. Therapeutic uses include treatment of viral infections of the brain or other diseased conditions where introduction of an antibody into the brain is required to treat the disease.

Brief Description of the Drawings

Figure 1 is a graph showing the increase in uptake of cationized IgG by brain capillaries at both 4°C and 37°C. The results are expressed as percent uptake of IgG per milligram of IgG.

Figure 2 is a graph which also shows the increase in uptake of cationized IgG by brain capillaries. The results are expressed as percent uptake of IgG versus amount of brain capillaries.

Figure 3 is a graph showing the uptake of cationized IgG plotted versus the concentration of either native IgG or cationized IgG.

Detailed Description of the Invention

The present invention involves the transport of antibodies through the brain capillary wall, i.e. the blood-brain barrier (BBB). The nature of the blood-brain barrier and problems associated with transport of peptides and proteins therethrough is set forth in "Receptor-Mediated Peptide Transport through the Blood-Brain Barrier" (W.M. Pardridge, *Endocrine Reviews*, Vol. 7, No. 3, August 1986, pages 314 -330), the contents of which is hereby incorporated by reference.

The present invention has wide application to any antibody which is useful in diagnosing or treating brain disorders. Antibodies in general do not readily cross the blood-brain barrier. This is due to the acidic or neutral character of antibodies. It was discovered that the uptake or transport of antibodies into the brain can be greatly increased by cationizing the antibodies to form cationized antibodies having an isoelectric point of between about 8.0 to 11.0.

Antibodies are proteins which have both positive and negative charges with the number of each depending upon the pH of the antibody solution.

The pH at which the positive and negative charges are equal is called the "isoelectric point" (pI). Techniques for measuring the pI of a given antibody or protein are well known and generally involve isoelectric focusing according to conventional electrophoresis procedure. As previously mentioned, most antibodies have an isoelectric point of between about 5 to 6.

The relatively low isoelectric point of antibodies is due to the presence of carboxyl groups on the surface of the antibodies. The present invention involves substituting basic groups in place of a sufficient number of surface carboxyl groups to increase the pI of the antibody to between about 8.0 to 11.0. Isoelectric points of between about 8.0 to 9.0 are preferred with isoelectric points of around 8.5 being especially preferred. The degree of cationization should be as high as possible without causing the antibody to form into aggregates. Higher pI's are preferred because the rate of transport of the antibody across the blood-brain barrier increases with increasing pI. However, this must be offset by the increasing possibility of antibody aggregate formation at higher levels of cationization.

Cationization of the antibody can be carried out according to any of the known procedures for displacing surface carboxyl groups on proteins with basic cations. Preferred cationization agents include amine compounds such as hexamethylenediamine and related amine compounds. Hexamethylenediamine is the preferred cationization agent because it is widely available and the techniques for its use in cationizing proteins are well known. The amount of cationizing agent and the conditions for reaction with the antibody can be varied so long as the final cationized antibody has a pI within the above-mentioned range required for blood-brain barrier transport.

The particular antibodies which can be used are virtually unlimited, provided that they have some diagnostic or therapeutic use in connection with the brain. Monoclonal antibodies are preferred because of their increased diagnostic or therapeutic potential. Typical antibodies which can be cationized for blood-brain barrier transcytosis are antibodies to one or more of the antigenic portions of peptides specific to Alzheimer's disease (Pardridge, W.M. et al., *Amyloid Angiopathy of Alzheimer's Disease: Amino Acid Composition and Partial Sequence of a 4,200 - Dalton Peptide Isolated from Cortical Microvessels*, *Journal of Neurochemistry*, 1987, pages 001 - 008). Antibodies to such specific peptides can be tagged with a radioactive tracer or other identifier and then cationized to a pI of 8.5 with hexamethylenediamine. The resulting tagged and cationized antibody can then be administered intravenously to the patient using a suitable pharmaceu-

3

5

EP 0 557 270 B1

6

tically acceptable carrier solution. The tagged and cationized antibody will cross the blood-brain barrier and enter the brain where it will bind to any of the peptides which are unique to Alzheimer's disease. Detection of the bond tagged and cationized antibody which is bound to the specific peptides is then performed by convention of neuroimaging techniques, such as external detection nuclide counting.

Other diagnostic antibodies which can be cationized to provide entry into the brain include antibodies for use in detecting various types of brain tumors. For example, monoclonal antibodies to tumor specific proteins such as glial fibrillary acidic protein (GFAP) can be prepared by conventional and well known techniques for monoclonal antibody preparation. Antibodies to human DR antigen and human immunodeficiency virus HIV antigen are other examples.

The resulting monoclonal antibodies are treated with hexamethylenediamine or other cationization agents to increase the pI of the antibody to between about 8.0 to 11.0. The antibody can be tagged with a radioactive tracer prior to or after the cationization process. The resulting tumor specific cationized and tagged antibody is then administered to the patient intravenously for transport across the blood-brain barrier and binding to any tumor specific antigen. Detection of bound antibody is again accomplished by convention radionuclide scanning techniques.

Cationized antibodies for use in treating viral diseases such as AIDS or other disorders of the brain can also be prepared as set forth above. Once an antibody (preferably monoclonal) is prepared for a specific neurotropic virus or other infectious agent, the antibody is cationized to increase its pI to between about 8.0 to 11.0. The antibody is then administered intramuscularly or intravenously to the patient. The antibody is typically administered as a solution of antibody in a suitable pharmaceutical carrier such as saline buffer. The doses of cationized antibody administered for either diagnostic or therapeutic purposes will parallel the dosage levels established for non-cationized antibodies. Typical dosages range from 0.01 mg to 1 mg for diagnostic purposes and from 1 mg to 100 mg for therapeutic purposes.

Preferred antibodies include chimeric human antibody molecules designed to have reduced antigenicity, such as those antibodies having mouse antigen-binding domains with human constant region domains. Such chimeric antibodies have been disclosed by S. L. Morrison et al. (Chimeric Human Antibody Molecules: Mouse Antigen-binding Domains with Human Constant Region Domains, Proc. Nat'l. Acad. Sci. USA, November 1984, Vol. 81, pages 6851 - 6855).

Although hexamethylenediamine is the preferred compound for use in cationizing antibodies, other cationizing agents are possible. For example, ethylene diamine, N,N-dimethyl-1,3-propanediamine, or polylysine may be used. Cationization is catalyzed by carboxyl activation using N-ethyl,N'(3-dimethyl-aminopropyl) carbodiimide hydrochloride (EDAC) using the method described by Hoare and Koshland (A Method for the Quantitative Modification and Estimation of Carboxylic Acid Groups in Proteins, 1967, J. Biol. Chem. 342:2447-2453).

In order to prevent reductions in the immunoreactivity of an antibody during cationization, it is preferred that the antibody be pre-bound to the antigen of interest prior to cationization. This pre-binding with antigen effectively blocks the immunoreactive sites on the antibody and prevents them from being cationized. After cationization is complete and the pI of the antibody has been raised to the desired level between about 8.0 to 11.0, the cationized antibody is then treated to unbind the antigen from the antibody. The unbinding is accomplished according to well known procedures where the antibody-antigen complex is treated with an acid to break the antibody-antigen bond. The antibody is then recovered by column chromatography or other conventional separation and recovery technique.

As an example of practice, bovine IgG was cationized and tested against native bovine IgG as follows:

One gram of bovine immunoglobulin G was dissolved in 10 ml of water followed by dialysis at 4°C overnight against water. To this was added slowly 67 ml of 2 M hexamethylenediamine while stirring, and the pH was kept at 7.8. Thirty minutes later, 1 g of EDAC was added and the pH was maintained at 7.8 and the solution was stirred at room temperature for 3-4 hours. The material was then dialyzed against 40 liters of Water overnight at 4°C followed by evaporation to dryness the following day.

The cationized antibody and native bovine antibody were then radiolabelled with ³H-sodium borohydride using standard methods which have been described previously by Partridge et al. (Absence of Albumin Receptor on Brain Capillaries In Vivo or In Vitro, 1985, Am. J. Physiol. 249:E264-E267; Chimeric Peptides as a Vehicle for Peptide Pharmaceutical Delivery through the Blood-Brain Barrier, 1987, Biochem. Biophys. Res. Commun. 146:307-315). Bovine brain capillaries were isolated from fresh bovine brain and used as in vitro model system of the blood-brain barrier as reviewed by Partridge, W.M. (Receptor-Mediated Peptide Transport through the Blood-Brain Barrier, 1986, Endocrine Reviews 7:314-330).

4

7

EP 0 557 270 B1

8

The results of the above tests with bovine IgG are set forth in Figures 1, 2 and 3. In Figure 1, the percent uptake per milligram protein of (¹²⁵I) cationized IgG or (¹²⁵I) native IgG at either 37 °C or 4 °C is plotted versus incubation time. The labeled cationized or native IgG was incubated with isolated bovine brain capillaries, which are used as an in vitro model system of blood-brain barrier transport. The results show that the cationization procedure increases the uptake of the IgG by nearly 50-fold, and that this is partially inhibited by cold temperatures. In Figure 2, the percent uptake of (¹²⁵I) cationized IgG or (¹²⁵I) native IgG is plotted versus the amount of bovine brain capillary protein content in micrograms per tube. There is approximately a 25-fold increase in the uptake of the IgG following cationization.

In Figure 3, the percent uptake of (¹²⁵I) cationized IgG per milligram protein of isolated bovine brain capillary is plotted versus the concentration of unlabeled cationized IgG or native IgG. The data show that the uptake of (¹²⁵I) cationized IgG is completely independent of concentration of native IgG through three log orders of magnitude in concentration. However, the presence of unlabeled cationized IgG stimulates the uptake in low concentrations and greatly depresses the uptake in high concentrations. The concentration of cationized IgG which causes 50% inhibition is approximately 2.5 mg/ml or approximately 15 μM cationized IgG.

Further examples of practice are:

A monoclonal antibody may be prepared against a synthetic peptide corresponding to the 4200 Dalton amyloid peptide of Alzheimer's disease amyloid angiopathy (see Pardridge et al, Amyloid angiopathy of Alzheimer's disease: amino acid composition and partial sequence of a 4,200-Dalton peptide isolated from cortical microvessels, 1987 J. Neurochem 49.) This amyloid is deposited on the brain side of the BBB and, thus, a monoclonal antibody to the amyloid peptide cannot be used as a neuroimaging device unless the monoclonal antibody is transportable through the BBB. The monoclonal antibody to the synthetic amyloid peptide may be cationized using hexamethylenediamine and EDAC, in the presence of saturated concentrations of synthetic amyloid peptide (to protect the active antigen binding sites on the antibody), to an isoelectric point of between 8 - 11. The cationized antibody may then be separated from the antigen by gel filtration in the presence of 0.1 M glycine (pH = 2.5). The high molecular weight peak containing the cationized antibody is then neutralized to pH = 7.4 and is now suitable for radiolabelling using standard radionuclides such as technetium 99m or iodine-¹²⁵I.

A monoclonal antibody to human GFAP may be prepared by isolating GFAP from human autopsy brain using standard techniques or by isolating recombinant human GFAP from either a bacterial or a eukaryotic expressing system. The monoclonal antibody to GFAP may then be cationized using hexamethylenediamine and EDAC in the presence of high concentrations of GFAP, followed by separation of cationized antibody from antigen as described above. The cationized monoclonal antibody to human GFAP may then be radiolabelled with technetium 99m or iodine-¹²⁵I or other conventional radionuclides. The final preparation is a radiolabelled antibody to GFAP that is transportable through the BBB and may be used as a neuroimaging device for early detection of brain glial tumors.

Another example is the preparation of mouse-human chimeric antibody directed against the human DR-antigen. This mouse-human chimeric antibody may be cationized using hexamethylenediamine and EDAC in the presence of saturating concentrations of recombinant DR-antigen followed by separation of cationized antibody from the free DR-antigen. The cationized human-mouse chimeric monoclonal antibody to the human DR-antigen may then be administered subcutaneously to subjects with demyelinating diseases, such as multiple sclerosis, that have an immune basis, and the pathogenesis of which may be ameliorated by the administration of antibody against the DR-antigen. For example, Sriram and Steinman (Anti I-A Antibody Suppresses Active Encephalomyelitis: Treatment Model for Diseases Linked to IR Genes, 1983, J. Exp. Med. 158:1362-1367) have provided evidence that immune linked demyelinating diseases may be treated by administration of antibody against the class II histocompatibility antigen.

Claims

1. A method for preparing an antibody having an increased transfer rate across the blood-brain barrier comprising the step of treating said antibody with an effective amount of a cationization agent to replace surface carboxyl groups on the antibody with a more basic group and thereby form a cationized antibody having an isoelectric point of between about 8.0 to 11.0.
2. A method according to claim 1 wherein cationized antibodies having an isoelectric point of between about 8.0 to 9.0 are formed.
3. A method according to claim 2 wherein cationized antibodies having an isoelectric

5

point of about 8.5 are formed.

4. A method according to any one of claims 1 to 3 wherein said cationization agent is an amine cationization agent.
5. A method according to claim 4 wherein said amine cationization agent is hexamethylenediamine.
6. A method according to any one of claims 1 to 5 wherein said antibody is a monoclonal antibody.
7. A method according to any one of claims 1 to 6 including the additional steps of:
 reacting said antibody with an antigen for said antibody to form an antibody having immunoreactive sites blocked by said antigen prior to cationization of said antibody; and
 treating the cationized antibody to remove said antigens therefrom to provide a cationized antibody having unblocked immunoreactive sites.
8. A method as claimed in any one of claims 1 to 7 characterised in that the antibody is tagged with a detectable tag.
9. A method as claimed in claim 8 characterised in that the tag is radioactive.
10. One or more antibodies formed by the method of any one of claims 1 to 7 for use in the manufacture of a composition for use in therapy or diagnosis.
11. One or more antibodies as claimed in claim 10 wherein said antibodies are selected from the group consisting of antibodies to Alzheimer's disease amyloid peptide, GFAP protein, DR-antigen and HIV antigen.
12. A diagnostic or neuropharmaceutical composition comprising one or more cationized antibodies having an isoelectric point of between about 8.0 and 11.0 and displaying increased rates of transfer across the blood-brain barrier compared to non-cationized antibodies.
13. A composition as claimed in claim 12 characterised in that the antibodies have an isoelectric point of between about 8.0 to 9.0.
14. A composition as claimed in claim 13 characterised in that the antibodies have an isoelectric point of about 8.5.

15. A composition as claimed in any one of claims 12 to 14 characterised in that it comprises antibodies to one or more selected antigens.
16. A composition as claimed in any one of claims 12 to 15 characterised in that it comprises a monoclonal antibody.
17. A composition as claimed in any one of claims 12 to 16 characterised in that the antibodies are tagged with a detectable tag.
18. A composition as claimed in claim 17 characterised in that the tag is radioactive.
19. The use of a composition as claimed in any one of claims 12 to 18 in the preparation of a medicament for use in a non-invasive method for introducing antibodies directly across the blood-brain barrier by transcytosis.
20. A composition as claimed in any one of claims 12 to 18 for use in diagnosis.
21. A composition as claimed in any one of Claims 12 to 18 for use in therapy.

Patentansprüche

1. Verfahren zur Herstellung eines Antikörpers erhöhter Transferrate durch die Blut-Hirn-Schranke durch Behandeln des Antikörpers mit einer wirksamen Menge eines Kationisierungsmittels zum Ersatz oberflächlicher Carboxylgruppen auf dem Antikörper durch eine stärker basische Gruppe unter Bildung eines kationisierten Antikörpers eines isoelektrischen Punkts zwischen etwa 8,0 und 11,0.
2. Verfahren nach Anspruch 1, wobei kationisierte Antikörper eines isoelektrischen Punkts zwischen 8,0 und 9,0 gebildet werden.
3. Verfahren nach Anspruch 2, wobei kationisierte Antikörper eines isoelektrischen Punkts von etwa 8,5 gebildet werden.
4. Verfahren nach einem der Ansprüche 1 bis 3, wobei es sich bei dem Kationisierungsmittel um ein Aminkationisierungsmittel handelt.
5. Verfahren nach Anspruch 4, wobei es sich bei dem Aminkationisierungsmittel um Hexamethylen-diamin handelt.
6. Verfahren nach einem der Ansprüche 1 bis 5, wobei es sich bei dem Antikörper um einen monoklonalen Antikörper handelt.

11

EP 0 557 270 B1

12

7. Verfahren nach einem der Ansprüche 1 bis 6 mit folgenden zusätzlichen Stufen: Umsetzen des Antikörpers mit einem Antigen für den betreffenden Antikörper vor der Kationisierung des Antikörpers zur Bildung eines Antikörpers mit durch das Antigen blockierten immunreaktiven Stellen und Behandeln des kationisierten Antikörpers zur Entfernung der Antigene von diesem zur Bereitstellung eines kationisierten Antikörpers mit nicht blockierten immunreaktiven Stellen.
8. Verfahren nach einem der Ansprüche 1 bis 7, dadurch gekennzeichnet, daß der Antikörper mit einer nachweisbaren Markierung markiert wird.
9. Verfahren nach Anspruch 8, dadurch gekennzeichnet, daß die Markierung radioaktiv ist.
10. Ein oder mehrere Antikörper, hergestellt nach dem Verfahren nach einem der Ansprüche 1 bis 7 zur Verwendung bei der Herstellung einer zu therapeutischen oder diagnostischen Zwecken verwendbaren Zubereitung.
11. Ein oder mehrere Antikörper nach Anspruch 10, wobei die Antikörper aus der Gruppe Antikörper gegen Amyloidpeptid der Alzheimer'schen Krankheit, GFAP-Protein, DR-Antigen und HIV-Antigen ausgewählt sind.
12. Diagnostische oder neuropharmazeutische Zubereitung mit einem oder mehreren kationisierten Antikörper(n) mit einem isoelektrischen Punkt zwischen etwa 8,0 und 11,0 und im Vergleich zu nicht kationisierten Antikörpern erhöhten Transferraten durch die Blut-Hirnschranke.
13. Zubereitung nach Anspruch 12, dadurch gekennzeichnet, daß die Antikörper einen isoelektrischen Punkt zwischen etwa 8,0 und 9,0 aufweisen.
14. Zubereitung nach Anspruch 13, dadurch gekennzeichnet, daß die Antikörper einen isoelektrischen Punkt von etwa 8,5 aufweisen.
15. Zubereitung nach einem der Ansprüche 12 bis 14, dadurch gekennzeichnet, daß sie Antikörper gegen ein oder mehrere ausgewählte(s) Antigen(e) umfaßt.
16. Zubereitung nach einem der Ansprüche 12 bis 15, dadurch gekennzeichnet, daß sie einen monoklonalen Antikörper umfaßt.
17. Zubereitung nach einem der Ansprüche 12 bis 16, dadurch gekennzeichnet, daß die Antikörper mit einer nachweisbaren Markierung markiert sind.
18. Zubereitung nach Anspruch 17, dadurch gekennzeichnet, daß die Markierung radioaktiv ist.
19. Verwendung einer Zubereitung nach einem der Ansprüche 12 bis 18 bei der Herstellung eines Medikaments zur Verwendung im Rahmen einer nicht invasiven Methode zur direkten Zufuhr von Antikörpern durch die Blut-Hirnschranke durch Transzytose.
20. Zubereitung nach einem der Ansprüche 12 bis 18 zur Verwendung bei der Diagnose.
21. Zubereitung nach einem der Ansprüche 12 bis 18 zur Verwendung bei der Therapie.

Revendications

1. Procédé de préparation d'un anticorps ayant une vitesse de transfert accrue à travers la barrière hémato-encéphalique, selon lequel on traite cet anticorps avec une quantité efficace d'un agent de cationisation pour remplacer les groupes carboxyle superficiels sur l'anticorps, par des groupes plus basiques et former un anticorps cationisé ayant un point isoélectrique d'environ 8,0 à 11,0.
2. Procédé selon la revendication 1, dans lequel on forme des anticorps cationisés ayant un point isoélectrique d'environ 8,0 à 9,0.
3. Procédé selon la revendication 2, dans lequel on forme des anticorps cationisés ayant un point isoélectrique d'environ 8,5.
4. Procédé selon l'une quelconque des revendications 1 à 3, dans lequel l'agent de cationisation est un agent de cationisation aminé.
5. Procédé selon la revendication 4, dans lequel l'agent de cationisation aminé est l'hexaméthylènediamine.
6. Procédé selon l'une quelconque des revendications 1 à 5, dans lequel l'anticorps est un anticorps monoclonal.
7. Procédé selon l'une quelconque des revendications 1 à 6, comprenant les étapes supplémentaires consistant à faire réagir cet anticorps avec un antigène

7

13

EP 0 557 270 B1

14

- ne correspondant à cet anticorps, pour former un anticorps dont les sites immunoréactifs sont bloqués par ledit antigène, avant cationisation de l'anticorps; et
à traiter l'anticorps cationisé pour éliminer les antigènes de celui-ci et procurer un anticorps cationisé comportant des sites immunoréactifs non bloqués.
8. Procédé selon l'une quelconque des revendications 1 à 7, caractérisé en ce que l'anticorps est marqué avec un marqueur détectable.
9. Procédé selon la revendication 8, caractérisé en ce que le marqueur est radioactif.
10. Un ou plusieurs anticorps formés selon le procédé de l'une quelconque des revendications 1 à 7, destinés à être employés pour la préparation d'une composition destinée à une utilisation thérapeutique ou diagnostique.
11. Un ou plusieurs anticorps selon la revendication 10, ces anticorps étant choisis parmi les anticorps dirigés contre le peptide amyloïde de la maladie d'Alzheimer, la protéine GFAP, l'antigène DR et l'antigène du virus HIV.
12. Composition diagnostique ou neuropharmaceutique, comprenant un ou plusieurs anticorps cationisés ayant un point isoélectrique d'environ 8,0 à 11,0, et donnant lieu à des vitesses accrues de transfert à travers la barrière hémato-encéphalique, par comparaison à des anticorps non cationisés.
13. Composition selon la revendication 12, caractérisée en ce que les anticorps ont un point isoélectrique d'environ 8,0 à 9,0.
14. Composition selon la revendication 13, caractérisée en ce que les anticorps ont un point isoélectrique d'environ 8,5.
15. Composition selon l'une quelconque des revendications 12 à 14, caractérisée en ce qu'elle comprend des anticorps dirigés contre au moins un antigène choisi.
16. Composition selon l'une quelconque des revendications 12 à 15, caractérisée en ce qu'elle comprend un anticorps monoclonal.
17. Composition selon l'une quelconque des revendications 12 à 16, caractérisée en ce que les anticorps sont marqués avec un marqueur détectable.
18. Composition selon la revendication 17, caractérisée en ce que le marqueur est radioactif.
19. Utilisation d'une composition selon l'une quelconque des revendications 12 à 18, pour la préparation d'un médicament destiné à un emploi non invasif, pour introduire des anticorps directement à travers la barrière hémato-encéphalique par transcytose.
20. Composition selon l'une quelconque des revendications 12 à 18, pour une utilisation diagnostique.
21. Composition selon l'une quelconque des revendications 12 à 18, pour une utilisation thérapeutique.

8

EP 0 557 270 B1

Fig. 2.





Fig. 1.



EP 0 557 270 B1

Fig. 3.

